

PRECISE DETERMINATION OF MUON SHOWER CONTENT FROM SHOWER UNIVERSALITY PROPERTY

A. Yushkov^{1*}, M. Ambrosio¹, C. Aramo¹, F. Guarino^{1,2}, D. D'Urso¹, L. Valore¹

¹INFN Sezione di Napoli, 80125, via Cintia, Napoli, Italia

²Università di Napoli "Federico II", 80125, via Cintia, Napoli, Italia

Abstract

It is shown, that highly accurate estimation of muon shower content can be performed on the basis of knowledge of only vertical depth of shower maximum $X_{\max}^V$ and total signal in ground detector. The estimate is almost independent on primary energy and particle type and on zenith angle. The study is performed for 21500 showers, generated with CORSIKA 6.204 from spectrum E^{-1} in the energy range $\log 10(E)$ [eV]=18.5–20 and uniformly in $\cos^2 \theta$ in zenith angle interval $\theta = 0^\circ - 65^\circ$ for QGSJET II/Fluka interaction models.

Muon shower content is the key parameter for studies of primary mass composition and test of hadronic interaction models. Using universality property of extensive air showers (EAS) development [1–6] we propose a simple and precise method to determine muon shower content from vertical depth of shower maximum and total signal (signal in water tanks or in scintillator detector). The study is performed for 21500 showers, generated with CORSIKA 6.204 [7] from spectrum E^{-1} in the energy range $\log 10(E)$ [eV]=18.5–20 (with different statistics in 3 energy bins 18.5–19.0, 19.0–19.5 and 19.5–20.0) and uniformly in $\cos^2 \theta$ in zenith angle interval $\theta = 0^\circ - 65^\circ$ for QGSJET II [8–10]/Fluka [11, 12] interaction models. Electromagnetic (EM) component thinning was set to 10^{-6} , the observation level was at 870 g/cm². All longitudinal showers characteristics and charged particles density were taken directly from CORSIKA output files. The expected signal in Auger-like tanks was calculated according to the procedure described in [13, 14] with the use of the same GEANT 4 lookup tables as in [14].

From different aspects of universality of shower development we will be interested only in the dependence of electromagnetic and muon signals on the distance of shower maximum to the ground. We will begin our consideration from Auger-like experimental setup and consider signal in water Cherenkov tanks at 1000 meters. In this case one deals with the situation, where muon contribution to the detector is much larger compared to the EM one due to higher energy losses of muons in tanks. The common way to express universality of electromagnetic signal is to plot it against slant distance to the ground (Fig. 1 and also Figure 1 in [4]), showing its quasi-independence on primary particle type. The muon signal¹ functional dependence on slant distance to the ground DG is also very similar for both proton and iron, but there is a shift in the normalization (Fig. 1). Since iron showers reach $X_{\max}$ earlier than showers from protons, comparison of set of showers from p and Fe at equal DG means the comparison between showers with different zenith angles, but at the same development stage. Passing to comparison of shower characteristics dependence on vertical distance to ground DG^V reveals a very interesting property (see Fig. 2): in this case the similarity of functional dependence of muon and EM signals on DG^V between p and Fe primaries is preserved, but now also EM signal normalizations are different. This happens because one confronts showers, which have the same vertical distance from $X_{\max}^V$ to the ground, but once again proton showers are more inclined than iron ones and their EM component attenuates more while reaching the ground from $X_{\max}^V$. The ratio $S_{\text{EM}}^{\text{Fe}}/S_{\text{EM}}^{\text{p}}$ turns out to be almost equal to the $S_{\mu}^{\text{Fe}}/S_{\mu}^{\text{p}}$ one and this allows to state the

*Corresponding author; e-mail: yushkov@na.infn.it

¹Muon signal includes only signal from muons, crossing the tank, signal from electromagnetic particles, originating from muon decays is included in the electromagnetic signal.

new shower universality property: *the ratio of the muon signal to the EM one S_μ/S_{EM} is the same for all showers, reaching the maximum at the same vertical depth X_{max}^v , independently on the primary particle type, primary energy and incident zenith angle* (at the least for the energy and angular ranges considered here). This property is illustrated in Fig. 3, where it is shown the dependence of S_μ/S_{EM} on X_{max}^v for p, O and Fe primaries in four different energy bins. The functional dependence between X_{max}^v and S_μ/S_{EM} turns out to be very simple and quasi-universal for all energies and primaries. The function in the form

$$X_{\text{max}}^v = A(S_\mu/S_{\text{EM}} + a)^b \quad (1)$$

fits well the data and the fit parameters are quite stable across entire energy range. Having in hand the functional dependence of X_{max}^v on (S_μ/S_{EM}) and using $S_{\text{tot}} = S_{\text{EM}} + S_\mu$ one easily gets the equation, which allows to obtain muon signal from shower vertical depth and total signal in tanks:

$$S_\mu^{\text{fit}} = \frac{S_{\text{tot}}}{1 + 1/((X_{\text{max}}^v/A)^{1/b} - a)}. \quad (2)$$

We calculated the difference between the Monte-Carlo (MC) simulated muon signal S_μ^{MC} and the muon signal, obtained from the fit S_μ^{fit} , an example distributions of this value are shown in Fig. 4. In Table 1 we give mean and RMS values of such distributions for various energy bins, obtained with the unique set of fit parameters $A = 538$, $b = -0.25$ and $a = -0.22$. It is seen, that the estimates of muon signals are unbiased with less than 1% deviation of mean reconstructed muon signal from the MC one for all 3 primaries and the RMS values are small: 8% for protons and around 5% for oxygen and iron. Certainly, application of specific coefficients for every energy bin or narrowing of zenith angle interval, or using of more sophisticated fit functions can even slightly improve the performance of the method, which anyhow is good in its simple and universal form. The described universal dependence of S_μ/S_{EM} on X_{max}^v holds true in the wide interval of distances and in Fig. 5 we show examples for the distances 200 and 1500 meters from the core, though for distances closer to the core the function in the form (1) does not describe accurately the data in the entire angular range $0^\circ - 65^\circ$ and it is needed or to split it in two parts or to apply more complex parametrization.

The same universality principle holds true in the case of detectors using scintillators and effectively measuring density of charged particles [15]. We have performed the reconstruction of muon densities using the dependence of the ratio (muon density D_μ)/(electron density D_e) on X_{max}^v in the form same to (1) (see Table 2 and Fig. 6). It is seen, that when muons and electrons equally contribute to the detector signal the shower fluctuations play more important role and the accuracy of parametrization is only within 15%, though the estimate is still unbiased.

Hence, the new universality property allows to obtain accurate estimates of the muon signal, which are almost independent on the primary particle type, primary energy and zenith angle for various types of ground detectors. Taking in consideration that the shower universality property was established for different interaction models [1–6], we expect that the proposed approach to muon content derivation is not specific only to QGSJET II/Fluka case.

The discovered simple shower universality property in respect to (muon signal/EM signal) ratio, giving access to the muon shower content, can open new possibilities as in solution of the global problems, such as derivation of primary mass composition and understanding of hadronic interactions properties, so in the number of more particular tasks (e.g. estimations of primary energy on the basis of pure electromagnetic signal, primary particle type independent corrections to the missing energy in experiments using fluorescent light etc.).

Acknowledgements

We are very grateful to Maximo Ave and Fabian Schmidt for kind permission to use their GEANT 4 lookup tables in our calculations of signal from different particles in Auger water tanks.




Figure 1: EM and muon signals from proton (red squares) and iron (blue crosses) in water Cherenkov tanks at 1000 m vs slant distance from shower maximum to the ground DG in $\log 10(E)[eV]=18.9-19.0$ energy bin




Figure 2: EM and muon signals from proton (red squares) and iron (blue crosses) in water Cherenkov tanks at 1000 m vs vertical distance from shower maximum to the ground DG^V in $\log 10(E)[eV]=18.9-19.0$ energy bin




Figure 3: Ratio of signals in water Cherenkov tanks S_μ/S_{EM} at 1000 m vs vertical depth of shower maximum X_{max}^v in four energy bins. Protons — red squares, oxygen — brown diamonds, iron — blue crosses




Figure 4: Distributions of relative difference between MC simulated muon signals in Cherenkov water tanks S_μ^{MC} and muon signals derived from the fit S_μ^{fit} at 1000 m. Protons — red line, oxygen — brown line, iron — blue line.




Figure 5: Ratio of signals in water Cherenkov tanks S_μ/S_{EM} at 200 m and 1500 m vs vertical depth of shower maximum X_{max}^v in $\log 10(E)[\text{eV}] = 18.9-19.0$ energy bin. Protons — red squares, oxygen — brown diamonds, iron — blue crosses

Table 1: Means and RMS of distributions of relative difference between MC simulated muon signals in Cherenkov water tanks S_μ^{MC} and muon signals derived from the fit S_μ^{fit} at 1000 m (see also Fig. 4) $(S_\mu^{\text{MC}} - S_\mu^{\text{fit}})/S_\mu^{\text{MC}}$, %, calculated with the unique set of parameters for all energy bins: $A = 538$, $b = -0.25$, $a = -0.22$

$\log 10(E)$ [eV]	proton		oxygen		iron	
	Mean	RMS	Mean	RMS	Mean	RMS
18.5 – 18.6	-0.1	7.4	-0.3	4.7	-0.8	4.2
18.6 – 18.7	-0.3	7.2	-0.2	5.0	-0.8	3.9
18.7 – 18.8	0.1	8.1	-0.2	4.9	-0.8	4.5
18.8 – 18.9	-0.1	8.3	-0.3	5.2	-0.5	4.2
18.9 – 19.0	0.5	8.3	-0.2	4.8	-0.5	4.3
19.0 – 19.1	-0.1	7.2	0.4	5.2	-0.3	4.2
19.1 – 19.2	0.4	8.5	0.2	5.1	-0.1	4.6
19.2 – 19.3	0.3	7.8	0.3	5.1	-0.1	3.9
19.3 – 19.4	0.2	7.9	0.0	5.2	0.2	4.5
19.4 – 19.5	0.6	8.0	0.1	5.4	-0.1	4.3
19.5 – 19.6	1.0	7.5	0.3	6.0	0.1	4.4
19.6 – 19.7	0.2	7.7	-0.0	5.1	0.0	4.7
19.7 – 19.8	0.7	8.1	0.3	4.9	-0.2	4.8
19.8 – 19.9	0.2	7.2	0.5	4.9	0.4	4.9
19.9 – 20.0	1.2	9.0	0.2	5.8	-0.1	4.9
18.5 - 20.0	0.3	7.9	0.1	5.1	-0.2	4.4




Figure 6: Top: ratio of muon density to the electron one at 1000 m vs vertical depth of shower maximum $X_{\text{max}}^{\text{v}}$ for two energy bins; bottom: distributions of relative difference between MC simulated muon density D_{μ}^{MC} and muon density derived from the fit D_{μ}^{fit} at 1000 m. The data are given for two energy bins, protons — red, oxygen — brown, iron — blue.

Table 2: Means and RMS of distributions of relative difference between MC simulated muon density D_μ^{MC} and muon density derived from the fit D_μ^{fit} at 1000 m (see also Fig. 6) $(D_\mu^{\text{MC}} - D_\mu^{\text{fit}})/D_\mu^{\text{MC}}$, %, calculated with the unique set of parameters for all energy bins $A = 475$, $b = -0.28$, $a = -0.09$

$\log 10(E)$ [eV]	proton		oxygen		iron	
	Mean	RMS	Mean	RMS	Mean	RMS
18.5 – 18.6	-0.3	15	-0.3	11	-1.8	10
18.6 – 18.7	-1.0	15	-0.1	11	-1.4	10
18.7 – 18.8	-0.0	16	-0.8	11	-0.9	10
18.8 – 18.9	-1.7	16	-0.2	11	-0.9	10
18.9 – 19.0	1.1	17	-0.8	12	-0.4	11
19.0 – 19.1	-0.2	15	0.6	12	-1.1	11
19.1 – 19.2	0.7	16	0.3	12	-0.5	11
19.2 – 19.3	0.3	15	0.5	13	-0.0	11
19.3 – 19.4	0.8	16	-0.9	12	0.4	11
19.4 – 19.5	1.3	17	0.1	13	0.3	12
19.5 – 19.6	1.3	17	1.5	13	0.3	12
19.6 – 19.7	-1.2	16	-0.4	13	0.7	12
19.7 – 19.8	0.9	17	-0.7	14	-0.8	12
19.8 – 19.9	0.2	16	-0.1	13	0.6	13
19.9 – 20.0	1.6	19	0.5	14	0.4	12
18.5 - 20.0	0.3	16	-0.1	12	-0.4	11

References

- [1] M. Giller, A. Kacperczyk, J. Malinowski, et al., *Similarity of extensive air showers with respect to the shower age*, J. Phys. **G31** (2005), 947–958.
- [2] F. Nerling, J. Bluemer, R. Engel, and M. Risse, *Description of Cherenkov light production in high-energy air showers*, Astropart. Phys. **24** (2006), 421–437.
- [3] D. Gora, R. Engel, D. Heck, et al., *Universal lateral distribution of energy deposit in air showers and its application to shower reconstruction*, Astropart. Phys. **24** (2006), 484–494.
- [4] Fabian Schmidt, Maximo Ave, Lorenzo Cazon, and Aaron S. Chou, *Applying Extensive Air Shower Universality to Ground Detector Data*, Proc. of the 30th ICRC (Merida), vol. 4, 2007, pp. 601–604.
- [5] P. Lipari, *The Concepts of ‘Age’ and ‘Universality’ in Cosmic Ray Showers*, Phys. Rev. **79** (2009), 063001.
- [6] S. Lafebvre, R. Engel, H. Falcke, et al., *Universality of electron-positron distributions in extensive air showers*, Astropart. Phys. **31** (2009), 243–254.
- [7] D. Heck, J. Knapp, J. N. Capdevielle, et al., *CORSIKA: a Monte Carlo code to simulate extensive air showers*, Forschungszentrum, Karlsruhe, 1998, **FZKA 6019**.
- [8] S. Ostapchenko, *QGSJET-II: towards reliable description of very high energy hadronic interactions*, Nucl. Phys. B (Proc. Suppl.) **151** (2006), 143–146.
- [9] ———, *Non-linear screening effects in high energy hadronic interactions*, Phys. Rev. **D74** (2006), 014026.
- [10] ———, *On the re-summation of enhanced Pomeron diagrams*, Phys. Lett. **B636** (2006), 40–45.
- [11] G. Battistoni, S. Muraro, P. R. Sala, et al., *The FLUKA code: Description and benchmarking*, AIP Conf. Proc. **896** (2007), 31–49.
- [12] A. Ferrari, P. R. Sala, A. Fasso, and J. Ranft, *FLUKA: A multi-particle transport code*, CERN-2005-010.
- [13] Pierre Billoir, *A sampling procedure to regenerate particles in a ground detector from a ‘thinned’ air shower simulation output*, Astropart. Phys. **30** (2008), 270–285.
- [14] Fabian Schmidt, Maximo Ave, Lorenzo Cazon, and Aaron S. Chou, *A Model-Independent Method of Determining Energy Scale and Muon Number in Cosmic Ray Surface Detectors*, Astropart. Phys. **29** (2008), 355–365.
- [15] M. Nagano, D. Heck, K. Shinozaki, et al., *Comparison of AGASA data with CORSIKA simulation*, Astropart. Phys. **13** (2000), 277–294.